

Proposal to search for a monochromatic component of solar axions using ^{57}Fe *

Shigetaka Moriyama ^a

^aDepartment of Physics, School of Science, University of Tokyo,
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

An experimental method is introduced to search for almost monochromatic solar axions. In this method, we can search for solar axions by exploiting the axion-quark coupling, not relying on the axion-photon coupling at all. A recent experimental result of Krčmar is presented.

1. Introduction

The most attractive solution of the strong CP problem is to introduce the Peccei-Quinn global symmetry which is spontaneously broken at energy scale f_a [1]. Although the original axion model has been experimentally excluded, variant “invisible” axion models are still viable. Such models are referred to as hadronic [2] and Dine-Fischler-Srednicki-Zhitnitskii [3] axions. At present, these “invisible” axions are constrained by laboratory searches and by astrophysical and cosmological arguments. The well-known window for f_a which escapes all the phenomenological constraints is 10^{10} – 10^{12} GeV. This window is frequently quoted because it is cosmologically interesting.

Besides this, there is another window around 10^6 GeV for the hadronic axions. This is usually called the hadronic axion window. A careful study [4] of the window revealed that f_a in the range 3×10^5 – 3×10^6 GeV cannot be excluded by the existing arguments, because most of them were based on the axion-photon coupling which is the least known parameter among those describing the low energy dynamics of the hadronic axions. In addition, it is argued that f_a around 10^6 GeV is desirable to explain the $ee\gamma\gamma + \cancel{E}_T$ event in the CDF experiment [5]. Furthermore, a recent paper indicates that axion might be a hot

dark matter if $f_a \sim 10^6$ GeV [6].

Although several authors [7,8] proposed experiments to search for the axions with f_a around 10^6 GeV, all of the experiments clearly rely on the axion-photon coupling both at the source and at the detector. The methods exploit the Primakoff effect; photons in the Sun are converted into axions, which are usually called the solar axions, and they are re-converted into X rays in a laboratory. The experiments to search for the solar axions by the Tokyo group [9], by the SOLAX collaboration [10], and by a pioneer [11] rely on the axion-photon coupling. The experiment [12], in which an emission line arising from the radiative decay of axions in the halo of our Galaxy is searched for, also proceeded through the axion-photon coupling. Thus there have been no experimental alternatives to test the hadronic axion window without relying on the axion-photon coupling.

This encouraged me to invent a new method to detect axions by exploiting axion-quark couplings, but not by relying on the axion-photon coupling.

2. ^{57}Fe in the Sun

Because of the axion coupling to nucleons, there is another component of solar axions. If some nuclides in the Sun have $M1$ transitions and are excited thermally, axion emission from nuclear deexcitation could be also possible. ^{57}Fe can be a suitable axion emitter for the following reasons: (i) ^{57}Fe has an $M1$ transition between the

*Proceedings of the Axion Workshop, University of Florida, Gainesville, Florida, USA, 13–15 March 1998, edited by P. Sikivie, to be published in Nucl. Phys. B Proc. Suppl.

first excited state and the ground state, (ii) the first excitation energy of ^{57}Fe is 14.4 keV, which is not too high compared with the temperature in the center of the Sun (~ 1.3 keV), and (iii) ^{57}Fe is one of the stable isotopes of iron, which is exceptionally abundant among heavy elements in the Sun. The natural abundance of ^{57}Fe is 2.2%. If the axion exists, therefore, strong emission of axions is expected from this nuclide.

These monochromatic axions would excite the same kind of nuclide in a laboratory because the axions are Doppler broadened due to thermal motion of the axion emitter in the Sun, and thus some axions have energy suitable to excite the nuclide. Fig. 1 shows the principle of the proposed experiment.

I propose to search for the axions by detecting this excitation. Since both the emission and absorption occurs via the axion-nucleon coupling but not via the axion-photon coupling, this method is free from the uncertainty of the axion-photon coupling. In addition, this method has merits that there is no need to tune the detector to a mass of the axions and that the sensitive mass region can be much higher than that of the proposed experiment [8], in which it is restricted by high pressure of buffer gas.

3. Expected event rate

The event rate is calculated in Ref. [13]. Since it depends on a parameter S which characterizes the flavor singlet coupling, which is not accurately determined, it is plotted as a function of S in Fig. 2. In the figure, the experimental values of S obtained recently [15] are included. The most recent experimental result gives $S = 0.39 \pm 0.11$ and discussions in Ref. [16] indicate $S \sim 0.5$. If the true value of S is around 0.4 as suggested, the expected event rate is restricted between 0.1 and 2.5 events $\text{kg}^{-1}\text{day}^{-1}$. Since there are both the upper bound and lower bound of the expected event rate, experiments which have adequate sensitivity will definitely determine whether the axion exists with f_a in the hadronic axion window.

4. Experimental method

We now turn to a discussion of experimental realities. After the excitation of the nuclei by the axion, there are two possible ways to relax. One is the emission of a γ ray with an energy of 14.4 keV. The other is the emission of an internal conversion electron with an energy of 7.3 keV and atomic radiations. Since the attenuation length of the γ ray is $20 \mu\text{m}$ and the range of the electron is $0.2 \mu\text{m}$ in iron, it is difficult to detect the γ rays or electrons outside the iron. In addition, detectors should have low energy threshold, low background, and high energy resolution. However, these difficulties are possibly overcome by using a bolometric technique with an absorber which contains ^{57}Fe -enriched iron. The technique has much advantages compared with other techniques in these respects. It is generally accepted that a sensitivity down to $0.1 \text{ events kg}^{-1}\text{day}^{-1}$ is reachable with a bolometer for dark matter search. If we can utilize this technique for the proposed experiment, it is possible to obtain a definite result as discussed above.

Recently, an experimental result based on this proposal was reported. Krčmar *et al.* [17] used a simple method to detect ^{57}Fe solar axions. They put a thin disk made of ^{57}Fe in front of a silicon lithium detector. A disk of natural iron with the same dimensions was used for background measurement. In this situation, they can detect only gamma rays which escape from the target. Since this is very rare, the total efficiency is extremely low. Their limit is that the axion mass is smaller than 745 eV. Although this is far beyond the hadronic axion window, I think this is a very interesting result.

5. Summary

In summary, a new scheme to detect almost monochromatic solar axions using resonant excitation of ^{57}Fe is introduced. ^{57}Fe is rich in the Sun and its first excitation energy is low enough to be excited thermally. Therefore, one can expect the nuclear deexcitation accompanied with the axion emission. Because of the Doppler effect associated with the thermal motion of ^{57}Fe in the




Figure 1. Principle of the proposed experiment.

Sun, a small portion of the axions from the nuclide can be absorbed by the same kind of nuclide in a laboratory. The nuclide is considered as a high sensitive detector of the axions. The excitation rate is expected to be order of $1 \text{ day}^{-1} \text{ kg}^{-1}$. Although it is difficult to detect the excitation outside the iron, it is detected efficiently by a bolometric technique with an absorber which contains ^{57}Fe -enriched iron. I am planning an experiment to search for the monochromatic axions from the Sun in this new scheme.

Acknowledgements

I want to thank Dr. S. Chang for his kindness and for discussion with him.

REFERENCES

1. R. D. Peccei and H. R. Quinn, Phys. Rev. Lett. **38**, 1440 (1977); Phys. Rev. D **16**, 1791 (1977).
2. J. E. Kim, Phys. Rev. Lett. **43**, 103 (1979); M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. B **166**, 493 (1980).
3. M. Dine, W. Fischler, and M. Srednicki, Phys. Lett. B **104**, 199 (1981); A. R. Zhitnitskii, Yad. Fiz. **31**, 497 (1980) [Sov. J. Nucl. Phys. **31**, 260 (1980)].
4. S. Chang and K. Choi, Phys. Lett. B **316**, 51 (1993).




Figure 2. Bound for the excitation rate. Hatched area is excluded by the argument of red-giants evolution and SN1987A [14]. Also shown is the experimental values of S that are measured recently [15].

5. J. Hisano, K. Tobe, and T. Yanagida, Phys. Rev. D **55** (1997) 411.
6. T. Moroi and H. Murayama, hep-ph/9804291.
7. P. Sikivie, Phys. Rev. Lett. **51**, 1415 (1983); E. A. Paschos and K. Zioutas, Phys. Lett. B **323**, 367 (1994).
8. K. van Bibber *et al.*, Phys. Rev. D **39**, 2089 (1989); F. Hoogeveen and R. G. Stuart, Phys. Lett. B **286**, 165 (1992).
9. S. Moriyama, M. Minowa, T. Namba, Y. Inoue, Y. Takasu, and A. Yamamoto, RESCEU-23/98, hep-ex/9805026, submitted to Phys. Lett. B; M. Minowa, these proceedings.
10. F. T. Avignone III *et al.*, astro-ph/9708008, submitted to Phys. Lett. B, and these proceedings.
11. D. M. Lazarus *et al.*, Phys. Rev. Lett. **69** (1992) 2333.
12. M. T. Ressell, Phys. Rev. D **44**, 3001 (1991).
13. S. Moriyama, Phys. Rev. Lett. **75** (1995) 3222.
14. W. C. Haxton and K. Y. Lee, Phys. Rev. Lett.

- 66**, 2557 (1991).
15. E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. D **54**, 6620 (1996); E143 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **75**, 25 (1995); E143 Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **74**, 346 (1995); Spin Muon Collaboration, D. Adams *et al.*, Phys. Lett. B **329**, 399 (1994); E142 Collaboration, P. L. Anthony *et al.*, Phys. Rev. Lett. **71**, 959 (1993); Spin Muon Collaboration, B. Adeva *et al.*, Phys. Lett. B **302**, 533 (1993).
 16. Spin Muon Collaboration, D. Adams *et al.*, Phys. Rev. D **56**, 5330 (1997); G. Altarelli, R. D. Ball, S. Forte, and G. Ridolfi, Nucl. Phys. B **496**, 337 (1997).
 17. M. Krčmar *et al.*, IRB-ZEF-98/01, nucl-ex/9801005.